

Quality of Post-Discharge Follow-Up in Patients Admitted to TJUH with Intermediate High Risk Pulmonary Embolism

D. Kramer, P. Kirupaharan, M. Flamenbaum, M. Scharf

Jane and Leonard Korman Respiratory Institute at Thomas Jefferson University, Philadelphia, PA

Problem Definition

- Our institution has developed a multidisciplinary pulmonary embolism response team (PERT) to help in the management of patients with intermediate high risk pulmonary embolism
- While the PERT focus is on acute inpatient management, it is important that PERT patients have close follow-up to monitor for known future complications including thromboembolic pulmonary hypertension (CTEPH) and chronic pulmonary embolism syndrome
- CTEPH, in particular, is a potentially treatable disease with significant morbidity if left untreated
- Therefore, an underappreciated and important aspect of the PERT team should be the coordinated follow-up and long term care of these patients
- We wanted to better understand how consistently our patients with intermediate high risk PE had pulmonary and/or vascular medicine (JATS) outpatient follow-up with repeat TTE performed at 3 months

Aims For Improvement

- To evaluate the rate of outpatient follow-up and evaluation after intermediate high risk PE to identify patients with higher risk of long term morbidity and mortality

Intervention

- We reviewed 72 patients based on initial date of PERT evaluation between the dates of January 1st, 2019 – December 31st, 2019
- We determined which patients with RV strain on echocardiography had appropriate outpatient follow-up and TTE within 3 months post-discharge

Measurement and Results

- Of 72 patients with PERT activation, only 48 had right ventricular (RV) strain by TTE
- Patients with CTA evidence of RV strain without corroborating evidence on TTE were excluded in our follow-up data as their risk of CTEPH was likely negligible




Fig. 1

Lessons Learned

- A significant percentage (33.3%) of PERTs are ordered based on CTA findings of RV strain that end up not being confirmed on echocardiography
- Follow-up with Pulmonary and/or JATS for the higher risk patients with RV strain on TTE was 60.8%, with 71.4% having repeat TTE performed at 3 months

Next Steps

- Determine if patients without follow-up had appointments made on discharge
- We would then tailor our intervention to the major driving factor prohibiting follow-up
 - No appointment vs. non-compliance would require different interventions




Fig. 2

1. PERT Coordinator with phone call to patient within a few days post-discharge
2. Pulmonary fellow messages office on Day 1 of PERT consult
3. Formal education to patient about expectations and CTEPH on day of discharge
4. Script for 3 month TTE handed to patient prior to discharge

References

1. Kim NH, Delcroix M, Jais X, et al. Chronic thromboembolic pulmonary hypertension. Eur Respir J 2019;53(1). 10.1183/13993003.01915-2018
2. Konstantinides SV, Meyer G, Becattini C, et al. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). Eur Heart J 2020;41(4):543-603. 10.1093/eurheartj/ehz405
3. Rivera-Lebron B, McDaniel M, Ahrar K, et al. Diagnosis, Treatment and Follow Up of Acute Pulmonary Embolism: Consensus Practice from the PERT Consortium. Clin Appl Thromb Hemost 2019;25:1076029619853037. 10.1177/1076029619853037